Sky Valley Treatment Resources

Recovery Center EvergreenHealth Monroe (360) 794-1405

- women's services inpatient, day treatment, outpatient, detox services, pregnant Medication-assisted treatment (TAM), assessment, residential
- Sea Mar Behavioral Health Monroe (360) 805-3122
- MAT, chemical dependency services, mental health services
- Conquer Addiction Monroe (360) 453-7715
- RAT, psychotropic medication management, counseling
- **Compass Health Monroe** (360) 419-7530
- of age, with complex behavioral health needs and their families. services and support to Medicaid-eligible individuals, up to 20 years Wraparound teams provide comprehensive behavioral health

2721-222 (778) 1 noitqO lssbl

MAT, outpatient services, behavioral health services referrals

Prevent Opioid Misuse

locking bottle, box, or cabinet.

or expired medications. Call , bətnewnu , bəsunu **Ased ə Ae**T

locations. med-project.org/ code to visit <u>www.</u> or scan the QR 1-844-633-7765

> less-addictive options for pain. Talk with your doctor about

for a short time. Get Narcan to doctor about using a low dose medication for pain, ask your lf you need opioid

counter and prescription-in a Fock your meds—over-the-



take home.

For a list of local treatment/recovery resources, Visit https://snohomishoverdoseprevention.com/ find-treatment-or-support/ or scan the QR code.



佐県

16

Never Use Alone: 1-877-696-1996



Suicide & Crisis Lifeline: or text 1-866-



Call or text 988



Teen Link: Call 833-6546



민하고

Help Lines and Treatment Resources

A COMMUNITY COMING TOGETHER TO STOP SNOHOMISH COUNTY'S OPIOID EPIDEMIC 200HOWI2HOVEROOSEPREVENTION

SKY VALLEY moo.noifneverdesobrevOdeimodon2.www

.dləd of ərəd ər'əW Looking for information? Ready for a change?

POCKET RESOURCE GUIDE

Signs & Symptoms of an Overdose

- Pale, clammy skin
- Not breathing/very shallow breathing
- Deep snoring or gurgling
- Unresponsive to name, shaking, etc.
- Slowed heartbeat/pulse
- Blue or ashy fingertips and lips
- Small pupils

If someone is making odd sounds while asleep, wake them. If they don't respond, shake them, call their name, or rub your knuckles over their breastbone.

Never let someone "sleep it off."

Responding to an Overdose

 Call or have someone call
The Good Samaritan Law protects you and the person who overdosed from drug possession charges.

Narcan lasts 30-90 minutes. It's possible for someone to slip back into an overdose after Narcan wears off. A medical center can continue treatment.

- 2. Give 1 dose of Narcan
- Open package and remove device. Place your thumb on the plunger and two fingers on the nozzle. Do not test or prime the device.
- Press the tip of the nozzle into either nostril until your fingers touch the bottom of

the person's nose.

• Press the plunger firmly to release the dose.

3. Wait 3 minutes. If they're unresponsive, give a second dose in the other nostril.

4. If they are not breathing and you have CPR training, start CPR.

5. If they are breathing, place them in the recovery position. Do not leave them alone unless absolutely necessary.

6. It's common to feel withdrawal symptoms and confusion after getting Narcan. The person may not remember what happened. Try to keep them calm until help arrives.

Where to get more Narcan

Find Narcan

near you, or have it confidentially mailed for free. Use the QR code to go <u>https://</u> stopoverdose.org.

From a pharmacy: No

prescription needed.



Washington has a Statewide Standing Order that acts as a universal prescription.

Private insurance: Ask your insurance company about coverage and co-pay.

Medicaid: Narcan is free, with no limits to requesting additional boxes.

For Current Drug Users

Never use alone. Carry Narcan. Tell anyone with you where to find your Narcan.

Use your substance slowly. Start with small amounts. Stop or take less if something feels off.

The Sound Pathways Syringe Service Program offers harm reduction supplies, 1:1 syringe exchange, referrals, and more. Scan the QR code to visit <u>https://www.facebook.com/syringeservices</u>.

